

FEED PROGRAM			All measurements are mL per gallon based on 2lb per gallon stock tank concentrate.										
MIXING ORDER ↓	CLONE	VEG 2-3 weeks recommended			FLOWER								
	Pre-Soak & Feed	W1	W2	W3	W1	W2	W3	W4	W5	W6	W7	W8	W9
Balance	pH up	Balance: Recommended for batch tank mixing and Dosatron. Do not use with Netaflex. Pro Balance: Recommended for advanced irrigation systems like Netaflex.											
Pro Balance													
Pro Core	19	19	19	19	19	19	19	19	19	19	*19	**Swap Fade	
Fade					*Swap Core with Fade W8-9 of flower. This is environment and cultivar dependent. If the cultivar shows resiliency, use Fade (19 mL) weeks 7-9.							19	19
Pro Grow		32	32	32									
Pro Bloom	32				32	32	32	32	32	32	32	32	32
Cleanse	1	2-5	2-5	2-5	2-5	2-5	2-5	2-5	2-5	2-5	2-5	2-5	2-5
EC	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
PPM 500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500	1500
PPM 700	2100	2100	2100	2100	2100	2100	2100	2100	2100	2100	2100	2100	2100
pH	5.6	5.5-5.8 (Coco/Rockwool) 5.9-6.2 (Peat based mediums)			5.8-6.2 (Coco/Rockwool)			6.0-6.4 (Peat based mediums)					6.0-6.4 (All)
**When switching to Fade verify pH.													

SPRAY PROGRAM			All measurements are mL per gallon					
APPLICATION FREQUENCY			VEG 2-3 weeks recommended			FLOWER		
			W1	W2	W3	W1	W2	W3
IPM	Preventative	2x Week	60-90	60-90	60-90	60-90	60-90	60-90
	Pressure	3x Week	90-120	90-120	90-120	90-120	90-120	90-120
Stack	Maximize	2x Week	7	7	7	7	7	7
Can be mixed together with IPM								

PRE-SOAK

Coco

- pH 5.5 - 5.8
- EC 3.5 - 4.0

Rockwool

- pH 5.0 - 5.5
- EC 2.5 - 3.0

FLUSH

RO + Cleanse	10 mL
EC	<0.1
PPM 500	<50
PPM 700	<70
pH	6.0 - 6.4

Coco
Last 3 days

Rockwool
Last day

PRO IRRIGATION STRATEGY

PRO DOSAGE GUIDELINES

IPM PROCEDURES

NOTE: Adjust feed chart according to weeks needed to complete a run. Strain dependent.